



Armstrong, M. EG., Kirichek, O., Cairns, B. J., Green, J., Reeves, G. K., Beral, V. (2016). Relationship of Height to Site-Specific Fracture Risk in Postmenopausal Women. *Journal of Bone and Mineral Research*, 31(4), 725-731. <https://doi.org/10.1002/jbmr.2742>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1002/jbmr.2742](https://doi.org/10.1002/jbmr.2742)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Wiley at <http://onlinelibrary.wiley.com/doi/10.1002/jbmr.2742/abstract>. Please refer to any applicable terms of use of the publisher.


University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

SUPPLEMENTARY INFORMATION


Supplemental Figure 1: Minimally adjusted relative risks and 99% CIs per 10cm increase in measured height for incident hospital admission for fracture at various sites and total fractures in post-menopausal women^{a b}



^a Minimally adjusted for age and stratified by study region

^b Mean values of measured height within self-reported categories used for trend calculation


Supplemental Figure 2: Relative risks and 99% CIs per 10cm increase in measured height for incident fractures at various sites, excluding the first two years of follow-up^{a b}



^a Adjusted for age, socio-economic status, BMI, strenuous activity, smoking, alcohol consumption, use of HRT, diabetes diagnosis, history of prior fracture and history of osteoporosis, and stratified by study region

^b Mean values of measured height within self-reported categories used for trend calculation

Supplemental Figure 3: Relative risks and 99% CIs per 10cm increase in measured height for incident fractures at various sites, restricted to women without missing data^{a b}



^a Adjusted for age, socio-economic status, BMI, strenuous activity, smoking, alcohol consumption, use of HRT, diabetes diagnosis, history of prior fracture and history of osteoporosis, and stratified by study region

^b Mean values of measured height within self-reported categories used for trend calculation